



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2015

ZBTB17 (MIZ1) Is Important for the Cardiac Stress Response and a Novel Candidate Gene for Cardiomyopathy and Heart Failure

Buyandelger, Byambajav ; Mansfield, Catherine ; Kostin, Sawa ; et al

Abstract: BACKGROUND Mutations in sarcomeric and cytoskeletal proteins are a major cause of hereditary cardiomyopathies, but our knowledge remains incomplete as to how the genetic defects execute their effects. METHODS AND RESULTS We used cysteine and glycine-rich protein 3, a known cardiomyopathy gene, in a yeast 2-hybrid screen and identified zinc-finger and BTB domain-containing protein 17 (ZBTB17) as a novel interacting partner. ZBTB17 is a transcription factor that contains the peak association signal (rs10927875) at the replicated 1p36 cardiomyopathy locus. ZBTB17 expression protected cardiac myocytes from apoptosis in vitro and in a mouse model with cardiac myocyte-specific deletion of Zbtb17, which develops cardiomyopathy and fibrosis after biomechanical stress. ZBTB17 also regulated cardiac myocyte hypertrophy in vitro and in vivo in a calcineurin-dependent manner. CONCLUSIONS We revealed new functions for ZBTB17 in the heart, a transcription factor that may play a role as a novel cardiomyopathy gene.

DOI: <https://doi.org/10.1161/CIRCGENETICS.113.000690>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-189916>

Journal Article

Published Version

Originally published at:

Buyandelger, Byambajav; Mansfield, Catherine; Kostin, Sawa; et al (2015). ZBTB17 (MIZ1) Is Important for the Cardiac Stress Response and a Novel Candidate Gene for Cardiomyopathy and Heart Failure. *Circulation. Cardiovascular Genetics*, 8(5):643-652.

DOI: <https://doi.org/10.1161/CIRCGENETICS.113.000690>

ZBTB17 (MIZ1) Is Important for the Cardiac Stress Response and a Novel Candidate Gene for Cardiomyopathy and Heart Failure

Byambajav Buyandelger, PhD; Catherine Mansfield, PhD; Sawa Kostin, PhD; Onjee Choi, PhD; Angharad M. Roberts, PhD; James S. Ware, PhD; Francesco Mazzarotto, PhD; Francesco Pesce, PhD; Rachel Buchan, PhD; Rivka L. Isaacson, PhD; Josée Vouffo, PhD; Sylvia Gunkel, PhD; Gudrun Knöll; Sara J. McSweeney, PhD; Heming Wei, PhD; Andreas Perrot, MD; Conny Pfeiffer†; Mohammad Reza Toliat, MD; Kristina Ilieva, MS; Ewelina Krysztofinska, MS; Marina M. López-Olañeta; Jesús M. Gómez-Salineró, MS; Albrecht Schmidt, MD; Keat-Eng Ng, PhD; Niels Teucher, MD; Ju Chen, PhD; Martin Teichmann, PhD; Martin Eilers, PhD; Wilhelm Haverkamp, MD; Vera Regitz-Zagrosek, PhD; Gerd Hasenfuss, MD; Thomas Braun, PhD; Dudley J. Pennell, MD; Ian Gould, PhD; Paul J.R. Barton, PhD; Enrique Lara-Pezzi, PhD; Sebastian Schäfer, PhD; Norbert Hübner, MD; Leanne E. Felkin, MD; Declan P. O'Regan, MD; Thomas Brand, PhD; Hendrik Milting, PhD; Peter Nürnberg, PhD; Michael D. Schneider, PhD; Sanjay Prasad, MD; Enrico Petretto, PhD; Ralph Knöll, MD, PhD

Background—Mutations in sarcomeric and cytoskeletal proteins are a major cause of hereditary cardiomyopathies, but our knowledge remains incomplete as to how the genetic defects execute their effects.

Methods and Results—We used cysteine and glycine-rich protein 3, a known cardiomyopathy gene, in a yeast 2-hybrid screen and identified zinc-finger and BTB domain-containing protein 17 (*ZBTB17*) as a novel interacting partner. *ZBTB17* is a transcription factor that contains the peak association signal (rs10927875) at the replicated 1p36 cardiomyopathy locus. *ZBTB17* expression protected cardiac myocytes from apoptosis in vitro and in a mouse model with cardiac myocyte-specific deletion of *Zbtb17*, which develops cardiomyopathy and fibrosis after biomechanical stress. *ZBTB17* also regulated cardiac myocyte hypertrophy in vitro and in vivo in a calcineurin-dependent manner.

Conclusions—We revealed new functions for *ZBTB17* in the heart, a transcription factor that may play a role as a novel cardiomyopathy gene. (*Circ Cardiovasc Genet.* 2015;8:643-652. DOI: 10.1161/CIRCGENETICS.113.000690.)

Key Words: cardiomyopathies ■ genetics ■ heart failure ■ models, animal ■ mutation

Mutations in >50 genes, mostly encoding sarcomeric and MZ disc proteins, cause hypertrophic cardiomyopathy (HCM) and dilated cardiomyopathy (DCM) that lead to heart failure.¹ Genetic variation in a Z disc gene can cause various forms of cardiomyopathies such as HCM or DCM perhaps because of pleiotropic effects on survival and hypertrophic pathways.² The muscle LIM protein (cysteine and glycine-rich protein 3 [CSRP3]) is a Z disc protein involved in cardiac mechanosensation that is important for myocyte-specific survival pathways through interactions with other Z disc proteins.³⁻⁵ CSRP3 mutations have been found in both HCM and

DCM patients,^{3,4,6} and we used protein-protein interaction studies to identify new CSRP3-interacting partners that may be important for cardiac myocyte survival or cardiomyopathy.

Clinical Perspective on p 652

Methods

All primers and antibodies used are listed in Tables I to III in the Data Supplement. Tissue culture experiments were performed using primary neonatal and adult rat cardiac myocytes, and animal experiments were performed using conditional knockout (cKO-Zbtb17), overexpressing transgenic (TG-ZBTB17) and double-transgenic

Received December 16, 2013; accepted July 2, 2015.

†Connie Pfeiffer passed away in 2013. We dedicate this work to her.

The Data Supplement is available at <http://circgenetics.ahajournals.org/lookup/suppl/doi:10.1161/CIRCGENETICS.113.000690/-DC1>.

Correspondence to Ralph Knöll, MD, PhD, AstraZeneca R&D Mölndal, R&D, Innovative Medicines and Early Development, Karolinska Institutet, Cardiovascular and Metabolic Diseases iMed Pepparedsleden 1, SE-431 83 Mölndal, Sweden and ICMC (Integrated Cardio Metabolic Centre), NOVUM, Level 7A, SE 141 57 Huddinge. E-mail: ralph.knoell@ki.se

© 2015 American Heart Association, Inc.

Circ Cardiovasc Genet is available at <http://circgenetics.ahajournals.org>

DOI: 10.1161/CIRCGENETICS.113.000690

(TG-ZBTB17/KO-Ppp3cb) protein phosphatase 3, catalytic subunit, β isozyme (calcineurin A β , Ppp3cb) animals.

The procedures in animal studies have been followed in accordance with institutional guidelines. Human studies have been approved by an institutional review committee, and that the subjects gave informed consent.

Yeast 2-hybrid, Western blot, quantitative real-time polymerase chain reaction, and other assays were performed as described elsewhere.^{3–5,7} Statistical evaluations were performed by nonparametric Mann–Whitney *U* tests and unpaired Student *t* tests (more details are available in the Data Supplement).

Results

CSRP3 Interacts With ZBTB17

Yeast 2-hybrid screens using CSRP3 as bait³ classified α -actinin and telethonin as CSRP3-interacting proteins that are important determinants of cardiac function^{5,8} and also identified ZBTB17 (Figure 1A). ZBTB17 is a member of the poxvirus and zinc-finger (POZ or BTB) domain/zinc-finger transcription factor family with no previously described role in the heart.⁹ We confirmed the CSRP3–ZBTB17 interaction by coimmunoprecipitation in vivo (Figure 1B), cross-linking of recombinant proteins (Figure 1C), and immunohistochemistry studies (Figure 1D; Figure I in the Data Supplement). These data identify ZBTB17 as a novel CSRP3-interacting protein.

Analysis of a Human ZBTB17 Variant (T106M)

ZBTB17 is encoded on human chromosome 1 at the replicated 1p36 cardiomyopathy locus.^{10–12} The peak DCM association at this locus (rs10927875; $P=1.3\times 10^{-7}$)¹² is located in intron 2 of *ZBTB17*, with further evidence of association of this single nucleotide polymorphism with DCM in Chinese population.^{12,13}

Considering rare polymorphisms (allele frequency <0.1%; population prevalence of cardiomyopathies $\approx 1:500$), we identified a p.T106M variant in the Exome Variant Server. The p.T106M variant could not be stably expressed in cardiac myocytes (Figure 2A), bound poorly to CSRP3 (Figure 2B), caused structural abnormalities to the POZ domain (Figure 2C and 2D), and may represent a rare haploinsufficient polymorphism. Together, these data point to *ZBTB17* as a possible disease gene at the 1p36 cardiomyopathy locus and suggest a role for a transcription factor in heart failure, as recently shown for left ventricular non compaction cardiomyopathy.¹⁴

Analysis of ZBTB17 Function In Vitro

To study function in vitro, we overexpressed *ZBTB17* in cardiac myocytes and found that it induced hypertrophy (Figure 3A and 3B; Figure IIA in the Data Supplement). CSRP3 and other CSRP3-interacting proteins have important roles in cardiac myocyte survival, and this was also observed for ZBTB17 (Figure 3C; Figure IIB in the Data Supplement). Calcineurin activity is of central importance for prohypertrophic signaling and myocyte survival,^{15,16} and in both the hypertrophy and the cell death experiments inhibition of calcineurin attenuated the effects of ZBTB17, suggesting an interaction (Figure 3A and 3C). Hence, ZBTB17 regulates cardiac hypertrophy and cell survival, a recognized

dual-property of genes that are of central importance for cardiac myocyte biology.^{15–18}

Analysis of ZBTB17 Function In Vivo

Mice with global deletion of *Zbtb17* die because of a gastrulation defect.¹⁹ Therefore, we crossed floxed *Zbtb17* mice²⁰ with the MLC2v Cre deleter line²¹ to test *Zbtb17* gene function in cardiac myocyte-specific cKO (Figure 3A–3D in the Data Supplement). cKO mice were viable and, in the absence of a spontaneous functional cardiac phenotype, we used the established model of transverse aortic constriction (TAC)⁵ to unveil potential gene effects. After TAC, the left ventricles of cKO mice became dilated and exhibited impaired contractile performance, which are the cardinal features of DCM (Table 1). When compared with controls, failing cKO hearts had higher apoptotic events (8.5-fold; $P<0.001$), increased activated caspase 3 (6.3-fold; $P<0.001$), and marked replacement fibrosis (Figure 4A; Figure III E and Table II in the Data Supplement). These data show that *Zbtb17* modulates biomechanical stress-induced apoptosis and interstitial fibrosis in vivo, which are important in the pathobiology of both HCM and DCM.

To begin to understand the molecular mechanisms by which ZBTB17 regulates cardiac myocyte survival, based on the functions of ZBTB17 in other tissues,¹⁹ we combined our microarray data with apoptosis-specific gene expression array analysis (quantitative real-time polymerase chain reaction based, $n=84$) in cKO.TAC and control.TAC mouse hearts. We used Gene Set Enrichment Analysis²² to test whether differentially expressed genes by genome-wide microarray analysis were significantly enriched in the set of differentially expressed genes detected by apoptosis-specific quantitative real-time polymerase chain reaction array analysis and found significant concordance for a set of 8 genes ($P<0.05$, false discovery rate, <1%; Table 2). Among these, the proapoptotic genes were consistently upregulated in cKO, whereas the antiapoptotic genes, *Nol3* and *Tnfrsf1a*, were downregulated in cKO hearts after biomechanical stress, supporting a role for *Zbtb17* in the regulation of apoptosis. Six of 8 differentially expressed genes were also detected by ChIP-chip analysis to have binding sites (Table 2). Hence ZBTB17 regulates a transcriptional program that protects cells against apoptosis.

In addition, ZBTB17 overexpression drives hypertrophy in vitro (Figure 3). To examine further the effects of *ZBTB17* on hypertrophy we overexpressed *ZBTB17* in mice using a cardiac myocyte-specific promoter²³ (Figure IV in the Data Supplement). Transgenic mice (TG-ZBTB17) exhibited spontaneous cardiac hypertrophy with enlarged cardiac myocytes and increased *Nppa* expression (Table I in the Data Supplement; Figure 4B and 4C). In vitro experiments suggested a link between ZBTB17 and calcineurin, and ChIP-chip data showed binding of ZBTB17 to the calcineurin A β (Ppp3cb) promoter that we confirmed by ChIP quantitative real-time polymerase chain reaction and reporter assays (Figure 4D and 4E). There was increased expression of Ppp3cb and *Nfatc2* in TG-ZBTB17 hearts and decreased expression in cKO (Figure 4F; Figure VA in the Data Supplement). We crossed *ZBTB17* transgenic mice into a Ppp3cb-deficient




Figure 1. **A**, Analysis of the interaction between cysteine and glycine-rich protein 3 (CSRP3) and zinc-finger and BTB domain-containing protein 17 (ZBTB17) by yeast 2-hybrid assays. No interaction occurs between proteins when the ZBTB17-POZ domain is deleted (lower right quadrant). A total of 10^6 clones were screened, and 650 positive clones were identified,³ one of which encoded the 5' sequence of *Zbtb17*. **B**, Immunoprecipitation of ZBTB17 from heart lysates results in the coprecipitation of CSRP3. Input indicates mouse heart extract, a *Csrp3* antibody (sc-30274; Santa Cruz Biotechnology) has been used for immunoprecipitation, and a well-characterized *Zbtb17* (Santa Cruz H-190; sc 22837) antibody has been used to detect the protein. No antibody or an unrelated anti-Kir2.1 antibody served as negative controls. **C**, Cross-linking experiments of recombinant proteins confirm binding between ZBTB17 and CSRP3. **Top**, With increasing concentration of cross-linking reagent, the individual components decrease, whereas a large complex emerges. The complex contains multiple copies of each protein reflecting the oligomeric nature of the individual components. **Bottom left**, Positive control showing cross-linking between CSRP3 and Telethonin (TCAP), a known interaction. **Bottom right**, Negative controls: we attempted to cross-link CSRP3 separately with 2 different proteins that have no documented association but are known to cross-link well to their binding proteins. These were small glutamine-rich tetratricopeptide repeat-containing protein alpha (SGTA) and HLA-B associated transcript 3 (BAT3). As expected, increasing cross-linking reagents showed no binding between CSRP3 and the negative control proteins, but only the usual CSRP3 oligomers indicated with stars. **D**, Colocalization of ZBTB17 and CSRP3 in the nuclei of human cardiac myocytes. Arrows indicate typical cardiac myocyte nuclei that are positive for both ZBTB17 and CSRP3. **Lower left**, Three-dimensional image of the boxed region (**left middle**). **Lower right**, Three-dimensional colocalized color intensity image of ZBTB17 with CSRP3: mean percentage of nuclear ZBTB17 colocalized with nuclear CSRP3=79.4 \pm 5.5%; mean Pearson coefficient of colocalized volumes=0.88 (n=4, \approx 125 nuclei per heart).




Figure 2. **A**, Analysis of wild-type (WT)- and T106M-zinc-finger and BTB domain-containing protein 17 (ZBTB17) proteins via adenoviral overexpression in neonatal rat cardiac myocytes. Although WT-ZBTB17 protein is readily detectable, ZBTB17:p.T106M was not and is most likely unstable in eukaryotic cells. **B**, Interaction of T106M-ZBTB17 with cysteine and glycine-rich protein 3 (CSRP3) as studied via cross-link interaction. CSRP3 interacts with WT-ZBTB17 (**left**), but interacts less well with T106M-ZBTB17 (**right**). **C**, NMR ^1H - ^{15}N HSQC spectrum of WT (red) and T106M (black) ZBTB17-POZ domain with pertinent residues assigned based on predicted chemical shift. The mutation causes significant structural disruption to the POZ domain. **D**, The available ZBTB17-POZ structure was used to model the T106M mutation. Fifty nanoseconds of simulation of the WT and the T106M structures that have been run as the full quadrameric structures as in the PDB 2Q81. Red indicates the native and yellow the mutant ZBTB17, spheres represent the T and M residues. It can be clearly seen that the structures of the α -6, where the mutation is located, have changed. Also the simulations reveal that there is a much higher level of structural mobility for the mutant case. The mutation causes significant structural disruption to the POZ domain, a finding consistent with our NMR data, arrowheads indicate position 106, arrows indicate a position far away).

background that prevented cardiac myocyte hypertrophy and decreased Nppa expression, confirming the interaction of ZBTB17 with the calcineurin pathway (Figure 4B; Table I in the Data Supplement).

Discussion

CSRP3 mutations cause heart failure in various animal models^{4,24} and have been identified in various patients affected by DCM^{3,25} or HCM.^{26–28} Although many cardiomyopathy and heart failure models have been developed, the molecular mechanisms that link sarcomeric and Z disc proteins to these phenotypes remain not well understood. Here, we show that CSRP3 interacts with ZBTB17 and provide the first detailed analysis of this transcription factor in the cardiovascular system. In particular, we demonstrate that *ZBTB17*

causes cardiac myocyte hypertrophy and is essential for cell survival. The extent of hypertrophy in cardiac myocytes is comparable with the effects seen with trophic signals including mitogenic serum in these postmitotic cells.^{29,30} To date, ZBTB17 is the only known transcription factor to interact with CSRP3 and to be expressed in cardiac myocytes.³¹ The effects of ZBTB17 on hypertrophy and survival are not only restricted to the in vitro situation but also observed in genetically altered animals under in vivo conditions. Interestingly, ZBTB17-overexpressing transgenic animals did not develop heart failure because of massive increase in apoptosis as frequently seen in other genetically altered mouse models.^{32–35}

The combination of ChIP-chip, whole-gene expression, and apoptosis array data clearly point to an important role of ZBTB17 in orchestrating a cardioprotective gene expression




Figure 3. **A**, Analysis of zinc-finger and BTB domain-containing protein 17 (ZBTB17) overexpression in neonatal rat cardiac myocytes (NRCMs) as determined by impedance measurements. Scatter dot plot represents cell index values after 24 h ($n=10$, experiments done in duplicate). Dotted lines indicate cell indices in calcineurin inhibition experiments (Cyclosporin A [CsA]: 0.2 μ mol/L; $n=4$; Mann-Whitney U test was used for the comparison of groups, $**P<0.01$). **B**, Representative images of NRCMs with and without ZBTB17 adenoviral gene transfer. **C**, Effects of ZBTB17 overexpression on doxorubicin (DOX, 1 μ mol/L)-induced cell death ($n=4$). Dotted lines indicate data from calcineurin inhibition experiments (CsA: 0.2 μ mol/L; $n=4$; Mann-Whitney U test was used for the comparison of groups, $*P<0.05$).

program. This is indicated by the fact that proapoptotic genes were consistently upregulated, whereas antiapoptotic genes were downregulated in cKO hearts after biomechanical stress (Table 3).

We also show that ZBTB17 targets various calcineurin/NFAT genes and activates this pathway. This notion is supported by our whole-gene expression analysis, ChIP-chip assays, analysis of the calcineurin/NFAT pathway activation, promoter assays, and by the significant loss of hypertrophy when TG-ZBTB17 animals are crossed into the calcineurin

$A\beta$ -deficient background (Figures 3 and 4D and 4E). These data are also supported by our finding of enhanced calcineurin $A\beta$ and *Nfatc2* mRNAs in hearts of ZBTB17 transgenic animals and decreased calcineurin $A\beta$ and *Nfatc2* mRNAs in hearts of *Zbtb17* cKO animals after TAC (Figure 4F; Figure VA in the Data Supplement).

CSRP3 translocates into the nucleus on TAC³⁶ and is essential for the adaptation of cardiac myocytes to biomechanical stress.³⁷ CSRP3, like telethonin, is primarily expressed in muscle tissues. By interacting with ZBTB17, CSRP3 plays a

Table 1. Echocardiography of cKO-Zbtb17 Animals

Genotype/ Intervention	BW, g	LVIDd, mm	LVIDs, mm	FS%	h/r	HW/BW, mg/g	LV _{vol} s	LV _{vol} d	EF%
Control.SHAM	31.90	3.77	2.06	45.61	0.54	4.26	14.40	61.39	45.62
Control.TAC	31.01	3.44*	1.65*	52.01*	0.63*	5.03*	7.86*	49.04*	52.04*
cKO.SHAM	32.04	3.71	2.05	45.71	0.56	4.07	15.30	59.79	45.73
cKO.TAC	32.55	3.74‡	2.00‡	46.46‡	0.57	4.82	13.22‡	59.69‡	46.46‡

Functional data obtained via echocardiography 4 weeks after sham (SHAM) and transverse aortic constriction (TAC) operations in *Zbtb17* conditional knockout (cKO) and control animals. After TAC, cKO hearts enlarge (LVIDd and LVIDs) and decrease in function (FS%; control=flox/flox, Cre⁻ and cKO=flox/flox, Cre⁺). BW indicates body weight; EF, ejection fraction; h/r, left ventricular wall thickness divided by left ventricular radius; HW, heart weight; LVIDd, left ventricular internal diameter in diastole; LVIDs, left ventricular internal diameter in systole; LV_{vol}d, left ventricular diastolic volume; and LV_{vol}s, left ventricular systolic volume.

**P*<0.05, for control.SHAM vs control.TAC.

‡*P*<0.05.

‡*P*<0.01 for control.TAC vs cKO.TAC; n=6 per group. Student *t* test was used for the comparison of groups.

role in the initiation of myocyte-specific survival pathways or mechanoptosis (mechanosensitive types of cell death⁵). This notion is supported by the fact that *Csrp3*-deficient hearts exhibit increased rates of apoptosis,³⁸ hence the interaction of CSRP3 with ZBTB17 can also be described as a form of Z disc transcriptional coupling.³⁹

Calcineurin/NFAT signaling is an important mediator of cardiac hypertrophy,^{40,41} and this pathway is also known to exert cardioprotective effects by avoiding apoptosis during ischemia reperfusion,¹⁵ which emphasizes the effects of ZBTB17 on cell survival.

CSRP3 directly interacts with calcineurin and is required for its activation,⁴² thus by interacting with ZBTB17; CSRP3 introduces an additional level of control, which ensures an intermediate level of calcineurin/NFAT activation (Figure 4G). For a long time, c-myc is known to be induced during TAC,⁴³ but its downstream effects in cardiac myocytes remained elusive. However, c-myc is known to inhibit ZBTB17⁴⁴ and therefore may have adverse effects on survival and hypertrophy in cardiac myocytes. Interestingly, CSRP3 interacts with the aminoterminal POZ domain, which is predicted to interfere with ZBTB17 tetramerization.

An additional layer of complexity is added by the fact that ZBTB17 can be phosphorylated at position S428 by the Akt kinase, which is another important mediator of hypertrophy.⁴⁵ This post-translational modification enables the interaction of ZBTB17 with 14-3-3 η and leads to its inactivation, thus avoiding prolonged activation of ZBTB17, which can also have adverse effects.⁴⁶

In the Exome Variant Server database, there are 10 T106M variants annotated out of at least 13006 alleles of European-American and Afro-American origin combined. The Exome Variant Server does not provide data on individual phenotypes, but the frequency for this variant is certainly low and a disease causing or at least modifying role cannot be excluded. Nevertheless, a 2-hit hypothesis whereby a structural perturbation combines with a defect in a ZBTB17-mediated cell survival pathway could also lead to heart failure.

As a result, we expect this variant to impair calcineurin-dependent hypertrophy and being unable to efficiently protect cardiac myocytes from apoptosis. However, this variant may also have specific effects on remodeling and

hypertrophy⁴⁷ by specifically affecting protein degradation pathways, for example, via the ubiquitin proteasome system,⁴⁸ and aside from affecting its interaction with CSRP3 (Figure 2B), may specifically interfere with other protein/protein interactions.

ZBTB17 is a transcription factor that contains the peak association signal (rs10927875) at the replicated 1p36 cardiomyopathy locus. Although other genes in the genomic region of *ZBTB17*, such as *HSPB7* and *CLCNKA*, have been suggested as possible candidate genes by genome-wide association study,^{10,49} *ZBTB17* might well be another novel DCM-associated gene. However, except for this report, the effects of ZBTB17 on the cardiovascular system have never been analyzed before. It is interesting to note that *HSPB7*, which encodes heat shock protein 27, a factor involved in the immediate stress response, *CLCNKA*, which encodes a K_a renal chloride channel involved in the regulation of hypo-osmotic cell stretch, as well as *ZBTB17*, are all involved in the cellular primary stress response and survival signaling.

At least 50 single genes have been identified as linked to familial DCM, some of which encode proteins of the sarcomere,⁵⁰ costamere, Z disc,^{3,4} and nuclear membrane,⁵¹ whereas others function as phosphatases and transcriptional activators (EYA4).^{1,52} Unfortunately, not much is known about how these effects are linked to changes in gene transcription. A recent study found profibrotic gene expression profiles in HCM mouse models and identified transforming growth factor beta 1 (*TGF β 1*) as the initiating event, but the underlying transcriptional events remained elusive.⁵³ However, ZBTB17 may well play a role in these genetic circuits. Defective Z disc-mediated survival signaling may also contribute to the DCM phenotype observed in patients carrying truncating Titin (*TTN*) mutations.⁵⁰

In summary, we identify *ZBTB17* as a candidate for a new cardiomyopathy gene, which may also be important for heart failure syndromes in general, and suggest that its primary function is to protect cardiac myocytes from apoptosis⁵ through modulation of both hypertrophic and cell death pathways (Figure 4G).

Acknowledgments

Dr Molkenstein is kindly acknowledged for providing the calcineurin A β Promotor-Luciferase plasmid and calcineurin A β




Figure 4. **A**, Representative micrographs of fibrosis staining of mouse hearts after sham (SHAM) or transverse aortic constriction (TAC) procedures and quantification of fibrosis (4 data points per animal, 1 representative experiment of 2 shown; Mann-Whitney test was used for the comparison of groups; $P < 0.05$). **B**, Analysis of cardiac myocyte cross-sectional area under spontaneous conditions in wild-type (WT)-zinc-finger and BTB domain-containing protein 17 (ZBTB17), conditional knockout (cKO)-Zbtb17, transgene (TG)-ZBTB17 (≈ 150 cardiac myocytes per animal; $n = 5$ per group, Mann-Whitney U test was used for the comparison of groups, $P < 0.05$, $**P < 0.01$), and KO-Ppp3cb, TG-ZBTB17/KO-Ppp3cb (≈ 150 cardiac myocytes per animal; $n = 2$ per group). **C**, Upregulation of the prohypertrophic biomarker Nppa in TG-ZBTB17 hearts ($n = 2$ per group, experiments done in duplicate). **D**, Chromatin immunoprecipitation microarray quantitative real-time polymerase chain reaction analysis of Zbtb17 binding to the Ppp3cb promoter in vivo (1 representative experiment of 2 shown, Mann-Whitney U test was used for the comparison of groups, $P < 0.05$). **E**, Luciferase reporter assay of ZBTB17 activation of the Ppp3cb promoter via transient transfection ($n = 3$ –4 per group, Mann-Whitney U test was used for the comparison of groups, $P < 0.05$). **F**, Increased *Ppp3cb* and *Nfatc2* mRNA expression levels in TG-ZBTB17 (1 representative experiment of 2 shown, Mann-Whitney U test was used for the comparison of groups, $P < 0.05$). **G**, ZBTB17 is important for the adaptation of cardiac myocytes to biomechanical stress. Loss of ZBTB17 is associated with an increase in cardiac myocyte apoptosis and heart failure, whereas overexpression leads to activation of the calcineurin/nuclear factor of activated T-cell (NFAT) pathway and cardiac myocyte hypertrophy. MLP indicates muscle LIM protein or CSRP3.

Table 2. Summarized Expression Profiles and Zbtb17 Binding Sites of Target Genes Involved in Cardiac Myocyte-Specific Programmed Cell Death in cKO Animals (n=6 Per Group)

Apoptosis RT ² Profiler PCR Array data (SA Biosciences)					Whole-transcript array data (Affymetrix Microarray)			GSEA	ChIP-Chip
Gene	Control TAC	cKO TAC	Fold Change	P Value	Control TAC	cKO TAC	Fold Change	Contribution to Gene Enrichment	FDR Value of Binding
Casp1	0.0034	0.006	1.76	0.028	0.0193	0.0386	2	+	0.00214
Casp2	0.0098	0.0215	2.2	0.002	0.0341	0.0431	1.26	+	0.02509
Casp3	0.003	0.0044	1.5	0.062	0.0223	0.0583	2.61	+	0.05001
Casp9	0.0112	0.0207	1.86	0.057	−0.0216	0.0107	2.5	+	0.01765
Traf3	0.0044	0.0071	1.61	0.008	−0.057	−0.0105	1.82	+	0.04486
Casp12	0.0008	0.0015	1.84	0.085	0.069	0.1771	2.57	...	0.02754
Nol3	0.0282	0.0198	0.7	0.008	0.1438	0.0985	0.68	...	NBS
Tnfrsf1a	0.0157	0.0092	0.59	0.054	0.1712	−0.0728	−0.43	...	NBS

Directional change of genes is consistent with the antiapoptotic role of endogenous Zbtb17. cKO indicates conditional knockout; GSEA, Gene Set Enrichment Analysis; PCR, polymerase chain reaction; NBS, no ZBTB17 binding site; TAC, transverse aortic constriction.

(Ppp3cb)-deficient animals. Dr Wright is thanked for providing the ZBTB17-POZ plasmid.

Sources of Funding

Dr Knöll is supported by Leducq (13CVD04), DFG Kn 448/9-1, /10-1, /10-2, Fritz Thyssen Stiftung, FP7-PEOPLE-2011-IRSES Proposal No 291834-Acronym: SarcoSi, and British Heart Foundation (BHF) grants PG/11/34/28793, RG 11/20/29266, SI/11/2/28875 and KI/AZ Integrated Cardio Metabolic Center (ICMC) grants. Dr Isaacson was supported by a Medical Research Council (MRC) New Investigator Research Grant. Dr Perrot was funded by Experimental and Clinical Research Center. Dr Schneider is the British Heart Foundation Simon Marks Chair in Regenerative Cardiology and supported by BHF grants RE/08/002 and SI/11/2/28875. Drs Barton, Ware, and Roberts are supported by grants from the MRC UK, the NMRC Singapore, the British Heart Foundation, the Wellcome Trust and the Leducq Foundation, Heart Research UK, The Academy of Medical Sciences and Arthritis Research UK. This project was also supported by the NIHR Cardiovascular Biomedical Research Unit of Royal Brompton and Harefield NHS Foundation Trust and Imperial College London. Drs Barton and Lara-Pezzi were supported by EU FP7 (CardioNet-ITN-289600).

Disclosures

None.

Appendix

From the National Heart and Lung Institute, Faculty of Medicine, Imperial Centre for Translational and Experimental Medicine (B.B., C.M., O.C., A.M.R., J.S.W., F.M., F.P., R.B., G.K., S.J.M., K.-E.N., E.L.-P., M.D.S., R.K.), KI/AZ Integrated Cardio Metabolic Center (ICMC), Department of Medicine, Karolinska Institute, Stockholm, Sweden (B.B., R.K.); Division of Molecular Biosciences, Department of Life Sciences, Faculty of Natural Sciences (R.L.I., K.I., E.K.), NIHR Cardiovascular Biomedical Research Unit at Royal Brompton & Harefield NHS Foundation Trust (D.J.P., P.J.R.B., L.E.F., S.P.), Department of Chemistry and Institute of Chemical Biology (I.G.), and Medical Research Council (MRC) Clinical Sciences Centre, Faculty of Medicine, Hammersmith Hospital (D.P. O'R., E.P.), Imperial College London, London, United Kingdom; Heart Centre, Faculty of Medicine, Georg-August University of Göttingen, Göttingen (B.B., J.V., S.G., G.K., C.P., A.S., N.T., G.H., R.K.); Max-Planck-Institute for Heart & Lung Research, Department of Cardiac Development and Remodeling, Bad Nauheim, Germany (S.K., T.B.); Charité-Universitätsmedizin Berlin, Department of Cardiology, Experimental and Clinical

Table 3. In Vivo Analysis of TUNEL-Positive Cells, Activated Caspase 3, and Collagen I

Animals	TUNEL (n/1000 Cardiomyocytes)		Activated Caspase 3 (n/1000 Cardiomyocytes)		Collagen I (% Tissue Area)	
	Control.TAC	cKO.TAC	Control.TAC	cKO.TAC	Control.TAC	cKO.TAC
1	1.04	11.35	3.53	17.73	5.89	10.71
2	1.73	12.55	3.86	21.81	6.63	12.53
3	1.49	9.41	4.01	19.55	7.22	16.24
4	2.38	15.44	3.37	19.92	5.14	11.25
5	0.95	13.87	3.88	25.77	7.73	15.02
6	1.72	8.08	1.85	23.19	6.85	13.09
Mean±SD	1.55±0.5	11.78±2.5*	3.42±0.73	21.32±2.63*	6.57±0.85	13.29±1.98*

cKO indicates conditional knockout; RT2, real-time-2 referring quantitative real-time PCR; TAC, transverse aortic constriction; and TUNEL, terminal deoxynucleotidyl transferase dUTP nick end labeling.

Student *t* tests were used for the comparison of groups, data=mean±SD; Control.TAC: flox/flox, Cre[−]; cKO.TAC: flox/flox, Cre⁺; n=6 per group. **P*<0.001.

Research Center (ECRC), a Joint Cooperation Between the Charité Medical Faculty and the Max-Delbrück Center for Molecular Medicine, Berlin, Germany (A.P.); Department of Cardiology at Campus Virchow-Klinikum (A.P., W.H.) and Berlin Institute of Gender in Medicine (V.R.-Z.), Charité-Universitätsmedizin Berlin, Berlin, Germany; Cologne Center for Genomics (M.R.T., P.N.) and Cologne Excellence Cluster on Cellular Stress Responses in Aging-Associated Diseases (CECAD) (P.N.), Center for Molecular Medicine Cologne (CMMC) (P.N.), University of Cologne, Cologne, Germany; Cardiovascular Development and Repair Department, Centro Nacional de Investigaciones Cardiovasculares (CNIC), Madrid, Spain (M.M. L.-O., J.M.G.-S., E.L.-P.); Department of Medicine, University of California at San Diego (UCSD) (J.C.); Institut Européen de Chimie et Biologie, INSERM U869, Bordeaux, France (M.T.); University of Bordeaux, IECB, ARNA Laboratory, Equipe Labellisée Contre le Cancer, F33600 Pessac, France (M.T.); Theodor Boveri Institute, Biocenter, University of Würzburg, Department of Physiological Chemistry II, Würzburg, Germany (M.E.); Ruhr-Universität Bochum, Herz & Diabeteszentrum NRW, Erich & Hanna Klessmann-Institut, Department of Cardiogenetics, Bad Oeynhausen, Germany (H.M.); Heart Science Centre, National Heart & Lung Institute, Imperial College London, Harefield Middlesex, United Kingdom (L.E.F., T.B.); Duke-NUS Graduate Medical School, Singapore, Singapore (H.W., E.P.); National Heart Centre, Singapore, Singapore (H.W.); Max-Delbrück-Center for Molecular Medicine (MDC), Berlin, Germany (S.S., N.H.); and DZHK (German Centre for Cardiovascular Research), Partner Site Berlin, Germany (S.S., N.H.); AstraZeneca R&D Mölndal, R&D, Innovative Medicines and Early Development, Karolinska Institutet, Cardiovascular and Metabolic Diseases iMed Pepparedsleden 1, SE-431 83 Mölndal, Sweden (R.K.).

References

- McNally EM, Golbus JR, Puckelwartz MJ. Genetic mutations and mechanisms in dilated cardiomyopathy. *J Clin Invest*. 2013;123:19–26. doi: 10.1172/JCI126262.
- Bos JM, Ackerman MJ. Z-disc genes in hypertrophic cardiomyopathy: stretching the cardiomyopathies? *J Am Coll Cardiol*. 2010;55:1136–1138. doi: 10.1016/j.jacc.2009.12.016.
- Knöll R, Hoshijima M, Hoffman HM, Person V, Lorenzen-Schmidt I, Bang ML, et al. The cardiac mechanical stretch sensor machinery involves a Z disc complex that is defective in a subset of human dilated cardiomyopathy. *Cell*. 2002;111:943–955.
- Knöll R, Kostin S, Klede S, Savvatis K, Klinge L, Stehle I, et al. A common MLP (muscle LIM protein) variant is associated with cardiomyopathy. *Circ Res*. 2010;106:695–704. doi: 10.1161/CIRCRESAHA.109.206243.
- Knöll R, Linke WA, Zou P, Micioc S, Kostin S, Buyandelger B, et al. Telethonin deficiency is associated with maladaptation to biomechanical stress in the mammalian heart. *Circ Res*. 2011;109:758–769. doi: 10.1161/CIRCRESAHA.111.245787.
- Geier C, Perrot A, Ozcelik C, Binner P, Counsell D, Hoffmann K, et al. Mutations in the human muscle LIM protein gene in families with hypertrophic cardiomyopathy. *Circulation*. 2003;107:1390–1395.
- Knöll R, Postel R, Wang J, Krätzner R, Hennecke G, Vacaru AM, et al. Laminin- α 4 and integrin-linked kinase mutations cause human cardiomyopathy via simultaneous defects in cardiomyocytes and endothelial cells. *Circulation*. 2007;116:515–525. doi: 10.1161/CIRCULATIONAHA.107.689984.
- Louis HA, Pino JD, Schmeichel KL, Pomiès P, Beckerle MC. Comparison of three members of the cysteine-rich protein family reveals functional conservation and divergent patterns of gene expression. *J Biol Chem*. 1997;272:27484–27491.
- Stead MA, Trinh CH, Garnett JA, Carr SB, Baron AJ, Edwards TA, et al. A beta-sheet interaction interface directs the tetramerisation of the Miz-1 POZ domain. *J Mol Biol*. 2007;373:820–826. doi: 10.1016/j.jmb.2007.08.026.
- Matkovich SJ, Van Booven DJ, Hinds A, Kang MY, Druley TE, Valiancia FL, et al. Cardiac signaling genes exhibit unexpected sequence diversity in sporadic cardiomyopathy, revealing HSPB7 polymorphisms associated with disease. *J Clin Invest*. 2010;120:280–289. doi: 10.1172/JCI39085.
- Stark K, Esslinger UB, Reinhard W, Petrov G, Winkler T, Komajda M, et al. Genetic association study identifies HSPB7 as a risk gene for idiopathic dilated cardiomyopathy. *PLoS Genet*. 2010;6:e1001167. doi: 10.1371/journal.pgen.1001167.
- Villard E, Perret C, Gary F, Proust C, Dilanian G, Hengstenberg C, et al; Cardiogenics Consortium. A genome-wide association study identifies two loci associated with heart failure due to dilated cardiomyopathy. *Eur Heart J*. 2011;32:1065–1076. doi: 10.1093/eurheartj/ehr105.
- Li X, Luo R, Mo X, Jiang R, Kong H, Hua W, et al. Polymorphism of ZBTB17 gene is associated with idiopathic dilated cardiomyopathy: a case control study in a Han Chinese population. *Eur J Med Res*. 2013;18:10. doi: 10.1186/2047-783X-18-10.
- Arndt AK, Schafer S, Drenckhahn JD, Sabeh MK, Plovie ER, Caliebe A, et al. Fine mapping of the 1p36 deletion syndrome identifies mutation of PRDM16 as a cause of cardiomyopathy. *Am J Hum Genet*. 2013;93:67–77. doi: 10.1016/j.ajhg.2013.05.015.
- Bueno OF, Lips DJ, Kaiser RA, Wilkins BJ, Dai YS, Glascock BJ, et al. Calcineurin Abeta gene targeting predisposes the myocardium to acute ischemia-induced apoptosis and dysfunction. *Circ Res*. 2004;94:91–99. doi: 10.1161/01.RES.0000107197.99679.77.
- Molkentin JD, Lu JR, Antos CL, Markham B, Richardson J, Robbins J, et al. A calcineurin-dependent transcriptional pathway for cardiac hypertrophy. *Cell*. 1998;93:215–228.
- Cook SA, Matsui T, Li L, Rosenzweig A. Transcriptional effects of chronic Akt activation in the heart. *J Biol Chem*. 2002;277:22528–22533. doi: 10.1074/jbc.M201462200.
- Ruppert C, Deiss K, Herrmann S, Vidal M, Oezkur M, Gorski A, et al. Interference with ERK(Thr188) phosphorylation impairs pathological but not physiological cardiac hypertrophy. *Proc Natl Acad Sci U S A*. 2013;110:7440–7445. doi: 10.1073/pnas.1221999110.
- Adhikary S, Peukert K, Karsunky H, Beuger V, Lutz W, Elsässer HP, et al. Miz1 is required for early embryonic development during gastrulation. *Mol Cell Biol*. 2003;23:7648–7657.
- Gebhardt A, Kosan C, Herkert B, Möry T, Lutz W, Eilers M, et al. Miz1 is required for hair follicle structure and hair morphogenesis. *J Cell Sci*. 2007;120(Pt 15):2586–2593. doi: 10.1242/jcs.007104.
- Hirota H, Chen J, Betz UA, Rajewsky K, Gu Y, Ross J Jr, et al. Loss of a gp130 cardiac muscle cell survival pathway is a critical event in the onset of heart failure during biomechanical stress. *Cell*. 1999;97:189–198.
- Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA, et al. Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide expression profiles. *Proc Natl Acad Sci U S A*. 2005;102:15545–15550. doi: 10.1073/pnas.0506580102.
- Achterberg A, Ackermann M, Adams J, Ahrens J, Andeen K, Atlee DW, et al; IceCube Collaboration. Limits on the high-energy gamma and neutrino fluxes from the SGR 1806-20 giant flare of 27 December 2004 with the AMANDA-II detector. *Phys Rev Lett*. 2006;97:221101.
- Arber S, Hunter JJ, Ross J Jr, Hongo M, Sansig G, Borg J, et al. MLP-deficient mice exhibit a disruption of cardiac cytoarchitectural organization, dilated cardiomyopathy, and heart failure. *Cell*. 1997;88:393–403.
- Mohapatra B, Jimenez S, Lin JH, Bowles KR, Coveler KJ, Marx JG, et al. Mutations in the muscle LIM protein and alpha-actinin-2 genes in dilated cardiomyopathy and endocardial fibroelastosis. *Mol Genet Metab*. 2003;80:207–215.
- Bos JM, Poley RN, Ny M, Tester DJ, Xu X, Vatta M, et al. Genotype-phenotype relationships involving hypertrophic cardiomyopathy-associated mutations in titin, muscle LIM protein, and telethonin. *Mol Genet Metab*. 2006;88:78–85. doi: 10.1016/j.ymgme.2005.10.008.
- Geier C, Gehmlich K, Ehler E, Hassfeld S, Perrot A, Hayess K, et al. Beyond the sarcomere: CSRP3 mutations cause hypertrophic cardiomyopathy. *Hum Mol Genet*. 2008;17:2753–2765. doi: 10.1093/hmg/ddn160.
- Newman B, Cescon D, Woo A, Rakowski H, Eriksson MJ, Sole M, et al. W4R variant in CSRP3 encoding muscle LIM protein in a patient with hypertrophic cardiomyopathy. *Mol Genet Metab*. 2005;84:374–375. doi: 10.1016/j.ymgme.2004.11.013.
- Sadoshima J, Aoki H, Izumo S. Angiotensin II and serum differentially regulate expression of cyclins, activity of cyclin-dependent kinases, and phosphorylation of retinoblastoma gene product in neonatal cardiac myocytes. *Circ Res*. 1997;80:228–241.
- Ueno H, Perryman MB, Roberts R, Schneider MD. Differentiation of cardiac myocytes after mitogen withdrawal exhibits three sequential states of the ventricular growth response. *J Cell Biol*. 1988;107:1911–1918.
- Buyandelger B, Ng KE, Micioc S, Piotrowska I, Gunkel S, Ku CH, et al. MLP (muscle LIM protein) as a stress sensor in the heart. *Pflugers Arch*. 2011;462:135–142. doi: 10.1007/s00424-011-0961-2.

32. Dorn GW II. Mitochondrial pruning by Nix and BNip3: an essential function for cardiac-expressed death factors. *J Cardiovasc Transl Res*. 2010;3:374–383. doi: 10.1007/s12265-010-9174-x.
33. Adams JW, Sakata Y, Davis MG, Sah VP, Wang Y, Liggett SB, et al. Enhanced Galpha signaling: a common pathway mediates cardiac hypertrophy and apoptotic heart failure. *Proc Natl Acad Sci U S A*. 1998;95:10140–10145.
34. Diwan A, Krenz M, Syed FM, Wansapura J, Ren X, Koesters AG, et al. Inhibition of ischemic cardiomyocyte apoptosis through targeted ablation of Bnip3 restrains postinfarction remodeling in mice. *J Clin Invest*. 2007;117:2825–2833. doi: 10.1172/JCI32490.
35. Yussman MG, Toyokawa T, Odley A, Lynch RA, Wu G, Colbert MC, et al. Mitochondrial death protein Nix is induced in cardiac hypertrophy and triggers apoptotic cardiomyopathy. *Nat Med*. 2002;8:725–730. doi: 10.1038/nm719.
36. Boateng SY, Belin RJ, Geenen DL, Margulies KB, Martin JL, Hoshijima M, et al. Cardiac dysfunction and heart failure are associated with abnormalities in the subcellular distribution and amounts of oligomeric muscle LIM protein. *Am J Physiol Heart Circ Physiol*. 2007;292:H259–H269. doi: 10.1152/ajpheart.00766.2006.
37. Boateng SY, Senyo SE, Qi L, Goldspink PH, Russell B. Myocyte remodeling in response to hypertrophic stimuli requires nucleocytoplasmic shuttling of muscle LIM protein. *J Mol Cell Cardiol*. 2009;47:426–435. doi: 10.1016/j.yjmcc.2009.04.006.
38. Heineke J, Wollert KC, Osinska H, Sargent MA, York AJ, Robbins J, et al. Calcineurin protects the heart in a murine model of dilated cardiomyopathy. *J Mol Cell Cardiol*. 2010;48:1080–1087. doi: 10.1016/j.yjmcc.2009.10.012.
39. Knöll R, Buyandelger B, Lab M. The sarcomeric Z-disc and Z-discopathies. *J Biomed Biotechnol*. 2011;2011:569628. doi: 10.1155/2011/569628.
40. Dorn GW II, Force T. Protein kinase cascades in the regulation of cardiac hypertrophy. *J Clin Invest*. 2005;115:527–537. doi: 10.1172/JCI24178.
41. Heineke J, Molkentin JD. Regulation of cardiac hypertrophy by intracellular signalling pathways. *Nat Rev Mol Cell Biol*. 2006;7:589–600. doi: 10.1038/nrm1983.
42. Heineke J, Ruetten H, Willenbockel C, Gross SC, Naguib M, Schaefer A, et al. Attenuation of cardiac remodeling after myocardial infarction by muscle LIM protein-calcineurin signaling at the sarcomeric Z-disc. *Proc Natl Acad Sci U S A*. 2005;102:1655–1660. doi: 10.1073/pnas.0405488102.
43. Izumo S, Nadal-Ginard B, Mahdavi V. Protooncogene induction and reprogramming of cardiac gene expression produced by pressure overload. *Proc Natl Acad Sci U S A*. 1988;85:339–343.
44. Wanzel M, Herold S, Eilers M. Transcriptional repression by Myc. *Trends Cell Biol*. 2003;13:146–150.
45. Shiojima I, Sato K, Izumiya Y, Schiekofer S, Ito M, Liao R, et al. Disruption of coordinated cardiac hypertrophy and angiogenesis contributes to the transition to heart failure. *J Clin Invest*. 2005;115:2108–2118. doi: 10.1172/JCI24682.
46. Wanzel M, Kleine-Kohlbrecher D, Herold S, Hock A, Berns K, Park J, et al. Akt and 14-3-3eta regulate Miz1 to control cell-cycle arrest after DNA damage. *Nat Cell Biol*. 2005;7:30–41. doi: 10.1038/ncb1202.
47. Knöll R, Iaccarino G, Tarone G, Hilfiker-Kleiner D, Bauersachs J, Leite-Moreira AF, et al. European Society of Cardiology. Towards a re-definition of 'cardiac hypertrophy' through a rational characterization of left ventricular phenotypes: a position paper of the Working Group 'Myocardial Function' of the ESC. *Eur J Heart Fail*. 2011;13:811–819. doi: 10.1093/eurjhf/hfr071.
48. Schlossarek S, Carrier L. The ubiquitin-proteasome system in cardiomyopathies. *Curr Opin Cardiol*. 2011;26:190–195. doi: 10.1097/HCO.0b013e32834598fe.
49. Cappola TP, Matkovich SJ, Wang W, van Booven D, Li M, Wang X, et al. Loss-of-function DNA sequence variant in the CLCNKA chloride channel implicates the cardio-renal axis in interindividual heart failure risk variation. *Proc Natl Acad Sci U S A*. 2011;108:2456–2461. doi: 10.1073/pnas.1017494108.
50. Herman DS, Lam L, Taylor MR, Wang L, Teekakirikul P, Christodoulou D, et al. Truncations of titin causing dilated cardiomyopathy. *N Engl J Med*. 2012;366:619–628. doi: 10.1056/NEJMoa1110186.
51. Fatkin D, MacRae C, Sasaki T, Wolff MR, Porcu M, Frenneaux M, et al. Missense mutations in the rod domain of the lamin A/c gene as causes of dilated cardiomyopathy and conduction-system disease. *N Engl J Med*. 1999;341:1715–1724.
52. Schönberger J, Wang L, Shin JT, Kim SD, Depreux FF, Zhu H, et al. Mutation in the transcriptional coactivator EYA4 causes dilated cardiomyopathy and sensorineural hearing loss. *Nat Genet*. 2005;37:418–422. doi: 10.1038/ng1527.
53. Teekakirikul P, Eminaga S, Toka O, Alcalai R, Wang L, Wakimoto H, et al. Cardiac fibrosis in mice with hypertrophic cardiomyopathy is mediated by non-myocyte proliferation and requires Tgf-β. *J Clin Invest*. 2010;120:3520–3529. doi: 10.1172/JCI42028.

CLINICAL PERSPECTIVE

Mutations in >50 genes, mostly encoding sarcomeric and Z disc proteins, are a major cause of hereditary cardiomyopathies. However, most likely this list is incomplete, and, probably most importantly, the underlying molecular disease-causing mechanisms remain not well defined. Here, we used cysteine and glycine-rich protein 3 (also known as muscle LIM protein), a known cardiomyopathy gene, and identified zinc finger and BTB domain-containing protein 17 (*ZBTB17*) as a novel interacting partner. *ZBTB17* is a transcription factor that contains the peak association signal (rs10927875) at the replicated 1p36 cardiomyopathy locus, identified in various recent genome-wide association studies. Loss of *Zbtb17* in cardiac myocytes causes apoptosis and heart failure, whereas overexpression of *ZBTB17* drives calcineurin/nuclear factor of activated T-cell (NFAT)–dependent cardiac myocyte hypertrophy. Here, we provide the first detailed analysis of *ZBTB17* in the cardiovascular system and provide evidence for *ZBTB17* being a novel cardiomyopathy candidate gene. *ZBTB17* functions might be exploited for the development of innovative therapies to treat heart failure, an otherwise lethal condition.